

Compositeness condition at the next-to-leading order in the Nambu-Jona-Lasinio model

Keiichi Akama

Department of Physics, Saitama Medical College, Kawakado, Moroyama, Saitama, 350-04, Japan

(February 1, 2008)

Abstract

We investigate the compositeness condition in four dimensions at the non-trivial order (next-to-leading order in $1/N$ with the number N of the colors), taking the Nambu-Jona-Lasinio model as an example. In spite of its non-renormalizability, the compositeness condition can be successfully derived and solved in compact forms. The resultant problems are discussed.

Compositeness may be one of the most important concepts in physics to disclose the fundamental structure of the nature. Needless to say, molecules, atoms, nuclei and hadrons are composites, and quarks, leptons, gauge bosons, and/or Higgs scalar also could be, as is supposed in some models [1] – [6] of the beyond-the-standard-model physics. Furthermore composite states play essential roles in various phenomena, such as superconductivity, nuclear excitation etc.. Nevertheless many of the dynamical systems are complicated and hard to be solved except for some idealistic cases. So far, extensive efforts have been devoted to this subject from both the particular and the general points of view.

Among them the “compositeness condition” [6] – [10] in the field theory could be a powerful clue, and should be investigated more extensively. This is the condition for a field in a model to be a composite of other fields, and expressed as

$$Z = 0, \tag{1}$$

where Z is the renormalization constant of the field. Under the condition (1), the composite field which is absent in the original bare-particle spectrum appears in the physical renormalized spectrum. As is shown in [8], a non-renormalizable four-fermion-interaction model is equivalent to a renormalizable Yukawa type model with the condition (1). Then, what is urgent for the four-fermion model is to work out the compositeness condition. In principle, it holds at any higher order. In fact, Haymaker and Cooper [9] considered the Gross-Neveu model [11] as a $Z = 0$ limit of the two-dimensional σ model, and investigated its higher-order effects. In four dimensions, however, the compositeness condition has been derived and solved only at the lowest order [8], which merely reproduces the well known results in the earlier naive approach [12]. A true nontrivial developments would be obtained by going beyond it. In recent years, people attempted to incorporate the higher order effects by using the ladder Schwinger-Dyson equation or the renormalization group equation [5], [6], [10], [13]. However, they are by nature incomplete because they take into account only particular classes of higher order diagrams. The more rigorous and promising way is to use the equivalent Yukawa model with the compositeness condition mentioned above. The

purpose of this paper is to make a true nontrivial developments by deriving and solving the compositeness condition at the nontrivial order (next-to-leading order in $1/N$ with the number N of the colors), and confirm the footing of the model. Remarkably enough, the infinite sum of divergent diagrams can successfully be summed up to give compact forms, and its solution is very simple.

We consider the Nambu-Jona-Lasinio (NJL) model in its simplest form :

$$\mathcal{L}_N = \bar{\psi}_0 i \not{\partial} \psi_0 + f_0 |\bar{\psi}_{0L} \psi_{0R}|^2, \quad (2)$$

where $\psi_0 = (\psi_{01}, \psi_{02}, \dots, \psi_{0N})$ is a bare color N -plet fermion, f_0 is a bare coupling constant, the subscripts “L” and “R” indicate chiralities, and the subscript “0” indicates that it is a bare quantity. This model has $U(1)_L \otimes U(1)_R$ chiral symmetry. The system is equivalent to that described by the linearized Lagrangian [14]

$$\mathcal{L}'_N = \bar{\psi}_0 i \not{\partial} \psi_0 + (\bar{\psi}_{0L} \phi_1 \psi_{0R} + \text{h.c.}) - \frac{1}{f_0} |\phi_1|^2, \quad (3)$$

where ϕ_1 is an auxiliary field. In order to absorb the divergences we rescale the fields as

$$\psi_0 = \sqrt{Z_\psi} \psi, \quad \phi_1 = \sqrt{X} \phi, \quad (4)$$

where Z_ψ and X are redundant constants, and ψ and ϕ are the physical (renormalized) fields. Then the Lagrangian (3) becomes

$$\mathcal{L}'_N = Z_\psi \bar{\psi} i \not{\partial} \psi + Z_\psi \sqrt{X} (\bar{\psi}_L \phi \psi_R + \text{h.c.}) - \frac{X}{f_0} |\phi|^2. \quad (5)$$

Now we compare this with the renormalizable Yukawa model for the elementary fermion ψ_0 and the elementary boson ϕ_0 with the following Lagrangian

$$\mathcal{L}_Y = \bar{\psi}_0 i \not{\partial} \psi_0 + g_0 (\bar{\psi}_{0L} \phi_0 \psi_{0R} + \text{h.c.}) + |\partial_\mu \phi_0|^2 - m_0^2 |\phi_0|^2 - \lambda_0 |\phi_0|^4 \quad (6)$$

where m_0 is the bare mass of ϕ_0 , and g_0 and λ_0 are bare coupling constants. To absorb the divergences of the quantum loop diagrams, we renormalize the fields, coupling constants and mass as

$$\psi_0 = \sqrt{Z_\psi}\psi, \quad \phi_0 = \sqrt{Z_\phi}\phi, \quad (7)$$

$$Z_\psi\sqrt{Z_\phi}g_0 = Z_g g, \quad Z_\phi^2\lambda_0 = Z_\lambda\lambda, \quad Z_\phi m_0^2 = m^2 + \delta m^2. \quad (8)$$

where ψ , ϕ , g , λ and m are the renormalized fields, coupling constants and mass, respectively, and Z_ψ , Z_ϕ , Z_g , Z_λ and δm^2 are the renormalization constants. Then the Lagrangian $\mathcal{L}_Y$ becomes

$$\begin{aligned} \mathcal{L}_Y = & Z_\psi \bar{\psi} i \not{\partial} \psi + Z_g g (\bar{\psi}_L \phi \psi_R + \text{h.c.}) \\ & + Z_\phi |\partial_\mu \phi|^2 - (m^2 + \delta m^2) |\phi|^2 - Z_\lambda \lambda |\phi|^4. \end{aligned} \quad (9)$$

Now we can see that the Lagrangian (9) of the Yukawa-type model coincides with the Lagrangian (5) of the NJL model, if

$$Z_\phi = Z_\lambda = 0, \quad (10)$$

$$Z_\psi \sqrt{X} = Z_g g, \quad (11)$$

$$X/f_0 = -m^2 - \delta m^2. \quad (12)$$

The conditions (11) and (12) connect the parameters of the NJL and the Yukawa models, and lead to so-called “the gap equation” of the NJL model. On the other hand, (10) is the “compositeness condition” which imposes relations among the parameters of the Yukawa model so that it reduces to the NJL model. In terms of the bare parameters, (10) implies the limit $g_0 \rightarrow \infty$ and $\lambda_0/g_0^4 \rightarrow 0$. Though some bare quantities diverge, renormalized ones are finite. Thus the non-renormalizable NJL model is entirely equivalent to the renormalizable Yukawa model (with the cutoff fixed) under the compositeness condition (10) [8]. Then what is urgent for the NJL model is to work out the compositeness condition. For this purpose, we need to calculate Z_ϕ and Z_λ in the Yukawa model and to solve $Z_\phi = Z_\lambda = 0$ for g and λ . In principle, we can perform it order by order to any higher order.

For an illustration, we start with the lowest order which reproduces (see below) the well known results in the earlier naive approach [12]. The divergences from the diagrams (a) and (b) in Fig. 1 are absorbed by taking

$$Z_\phi = 1 - g^2 NI, \quad Z_\lambda \lambda = \lambda - g^4 NI, \quad (13)$$

where $I = \ln \Lambda^2 / 16\pi^2$ in the Pauli-Villars regularization, or $I = 1/8\pi^2(4-d)$ in the d -dimensional regularization. The compositeness condition $Z_\phi = Z_\lambda = 0$ with (13) is easily solved to give

$$g^2 = \lambda = 1/NI. \quad (14)$$

If $m^2 < 0$, then ϕ acquires a non-vanishing vacuum expectation value $\langle \phi \rangle$ and the chiral symmetry is broken, so that $\pi = \text{Im}\phi$ becomes the Nambu-Goldstone boson, and ψ and $\sigma = \text{Re}\phi - \langle \phi \rangle$ respectively acquire the masses

$$m_\psi = \langle \phi \rangle / \sqrt{NI}, \quad m_\sigma = 2\langle \phi \rangle / \sqrt{NI}. \quad (15)$$

The relations (14) and (15) reproduce the well known results of the original NJL model [12] with the chain approximation. In this sense the method of the compositeness condition is trivial at this order.

Now we turn to the non-trivial order, which is the purpose of this paper. First we note the fact that a ψ -loop insertion to a ϕ -line or a ϕ -vertex contributes a factor $O(g^2 NI) \approx O(1)$ (because of (14)), and does not change the order of magnitude. Then infinitely many higher order diagrams have the same order of magnitude and the naive expansion in g and λ fail. A proper expansion is that in $1/N$ and $1/I$ with the assignment $g^2 = O(1/NI)$ and $\lambda = O(1/NI)$. Here we retain the next-to-leading order in $1/N$ and the leading order in $1/I$. They consist of the infinitely many diagrams (c)–(g) in Fig. 1, together with their counter terms for the sub-diagram divergences.

Now we derive the contribution to Z_ϕ from the diagram (c) in Fig. 1. Each ψ -loop inserted to the internal ϕ -line with the momentum p contributes (in d -dimensional regularization)

$$g^2 NI c_1 (-p^2)^{1-\epsilon} \quad (16)$$

with $\epsilon = (4-d)/2$ and $c_1 \rightarrow 1$ as $d \rightarrow 4$. If l of ψ -loops (16) are inserted to the ϕ -line, the self-energy part of ψ with the momentum k is calculated to be

$$(g^2 I)^{l+1} N^l c_2 k / (-k^2)^{\epsilon(l+1)} + O(m^2) \quad (17)$$

with $c_2 \rightarrow 1$ as $d \rightarrow 4$. Then the diagram (c) in Fig. 1 contribute to Z_ϕ the term

$$(-g^2 I)^{l+2} N^{l+1} / (l+1)(l+2). \quad (18)$$

If we take into account the counter terms for the divergence of (16) and (17),

$$(g^2 I)^{l+2} N^{l+1} \sum_{k=0}^l \binom{l}{k} \left(\frac{(-1)^k}{(k+1)(k+2)} - \frac{(-1)^k}{(k+1)} \right) = -\frac{(g^2 I)^{l+2} N^{l+1}}{(l+1)(l+2)}. \quad (19)$$

Collecting all together, we finally obtain the expression

$$Z_\phi = 1 - g^2 N I - g^2 I - \frac{1}{N} (1 - g^2 N I) \log(1 - g^2 N I) + O\left(\frac{1}{N^2}\right), \quad (20)$$

where the logarithmic term appear through the infinite sum of (19) over l . We can derive entirely the same expression as (20) also in the Pauli-Villars regularization.

Similarly the diagrams (d)–(g) in Fig. 1 together with their counter terms for the sub-diagram divergences contribute to $Z_\lambda \lambda$ the terms

$$\begin{aligned} (d) \quad & -g^4 I (g^2 N I)^l / l(l+1), \quad (e) \quad 10\lambda^2 I (g^2 N I)^l / (l+1), \\ (f) \quad & -20\lambda g^2 I (g^2 N I)^l / (l+1), \quad (g) \quad 10g^4 I (g^2 N I)^l / (l+1), \end{aligned} \quad (21)$$

where l is the number of the ψ -loops in the diagram. Collecting all together, we obtain the expression

$$\begin{aligned} Z_\lambda \lambda = & \lambda - g^4 N I + 8g^4 I + \frac{20(\lambda - g^2)^2 I}{1 - g^2 N I} \\ & - \frac{1}{N} \left[2g^2(1 - g^2 N I) + 20(\lambda - g^2) \right] \log(1 - g^2 N I) + O\left(\frac{1}{N^3}\right). \end{aligned} \quad (22)$$

The compositeness condition is obtained by putting $Z_\phi = Z_\lambda = 0$ in (20) and (22). Though it looks complicated at first sight, they are successfully solved giving the very simple solution

$$g^2 = \frac{1}{N I} \left[1 - \frac{1}{N} + O\left(\frac{1}{N^2}\right) \right], \quad \lambda = \frac{1}{N I} \left[1 - \frac{10}{N} + O\left(\frac{1}{N^2}\right) \right]. \quad (23)$$

If the chiral symmetry is spontaneously broken the fermion ψ and the scalar σ acquire masses

$$m_\psi = \frac{\langle\phi\rangle}{\sqrt{NI}} \left[1 - \frac{1}{2N} + O\left(\frac{1}{N^2}\right) \right], \quad m_\sigma = \frac{2\langle\phi\rangle}{\sqrt{NI}} \left[1 - \frac{5}{N} + O\left(\frac{1}{N^2}\right) \right]. \quad (24)$$

This means that the famous ratio $m_\sigma/m_\psi = 2$ in the lowest-order NJL model should be corrected as

$$\frac{m_\sigma}{m_\psi} = 2 \left[1 - \frac{9}{2N} + O\left(\frac{1}{N^2}\right) \right]. \quad (25)$$

The results are somewhat unpleasant for small N , in particular, for $N=3$ of our practical interest. The next to leading contributions to Z_λ , and accordingly, those to λ are too large. Furthermore λ is negative, and hence the Higgs potential becomes unstable. Its true stability may depend on the effective potential at large ϕ . The difference, however, comes from the convergent diagrams, and is suppressed by a factor of $1/I$ compared with the Higgs potential itself. The non-leading effects in $1/I$ may improve the situation. A possible way to go around the difficulty may be to take into account the gauge interactions and/or higher-dimensional interactions [15]. On the other hand, it may be too simple-minded to take the cutoffs of ψ and ϕ as equal. Introduction of the two cutoff scales could improve the situation [16]. These possible improvements are now under consideration.

We have derived and solved the compositeness condition at the non-trivial order, which has turned out to have large contributions at $N = 3$. It is straightforward to extend the investigation to more realistic cases of, for example, $SU(2)_L \otimes SU(2)_R$ for hadron physics, or $SU(2)_L \otimes U(1)_Y$ for electroweak theory. In the realistic model, we should take into account the effects of gauge bosons also. The present method is also applicable to the models of composite fermions and composite gauge bosons [17], and to the induced gravity theory (pregeometry) [18]. Another challenge may be to apply it to the QCD bound states. Recently some people used the compositeness condition as a boundary condition of the renormalization group equation [6]. It may also require the higher order consideration as is given here. More thorough investigations including further higher orders would lead to deeper insights into the compositeness.

The author would like to thank Professors H. Terazawa, A. J. Macfarlane, R. W. Haymaker, I. Ichinose, M. Suzuki, T. Hattori, T. Matsuki, H. Murayama, H. Nakano, M. Yasuè,

and Y. Watabiki for stimulating and critical discussions.


REFERENCES

- [1] J.C. Pati and A. Salam, Phys. Rev. **D10** (1974) 275.
- [2] K. Akama and H. Terazawa, INS-Rep-257 (1976); H. Terazawa, Y. Chikashige and K. Akama, Phys. Rev. **D15** (1977) 480.
- [3] G. 't Hooft, in *Recent Developments in Gauge Theories*, Cargèse, 1979, edited by G. 't Hooft *et al.* (Plenum Press, New York, 1980), p.135.
- [4] L. Susskind, Phys. Rev. **D19** (1979) 2691.
- [5] V.A. Miransky, M. Tanabashi and K. Yamawaki, Phys. Lett. **B221** (1989) 177; Mod. Phys. Lett. **A4** (1989) 1043.
- [6] W.A. Bardeen, C.T. Hill and M. Lindner, Phys. Rev. **D41** (1990) 1647.
- [7] B. Jouvett, Nuovo Cim. **5** (1956) 1133; M. T. Vaughn, R. Aaron and R. D. Amado, Phys. Rev. **124** (1961) 1258; A. Salam, Nuovo Cim. **25** (1962) 224; S. Weinberg, Phys. Rev. **130** (1963) 776.
- [8] D. Lurie and A. J. Macfarlane, Phys. Rev. **136** (1964) B816; T. Eguchi, Phys. Rev. **D14** (1976) 2755; **D17** (1978) 611; D. Campbell, F. Cooper, G. S. Guralnik and N. Snyderman, Phys. Rev. **D19** (1979) 549; K. Shizuya, Phys. Rev. **D21** (1980) 2327.
- [9] R. W. Haymaker and F. Cooper, Phys. Rev. **D19** (1979) 562.
- [10] F. Cooper and J. Perez-Mercader, Phys. Rev. **D43** (1991) 4129; D. Lurie and G.B. Tupper, Phys. Rev. **D47** (1993) 3580.
- [11] D. J. Gross and A. Neveu, Phys. Rev. **D10** (1974) 3235.
- [12] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345.
- [13] J. Zinn-Justin, Nucl. Phys. **B367** (1991) 105; H. Yamamoto and I. Ichinose, Nucl. Phys. **B370** (1992) 695; P. Fishbane, R.E. Norton and T.N. Truong, Phys. Rev. **D46** (1992)

- 1768; P. Fishbane and R.E. Norton, Phys. Rev. **D48** (1993) 4924; R.S. Willey, Phys. Rev. **D48** (1993) 2877; J. A. Gracey, Phys. Lett. **B308** (1993) 65; **B342** (1995) 297.
- [14] T. Kugo, Prog. Theor. Phys. **55** (1976) 2032; T. Kikkawa, Prog. Theor. Phys. **56** (1976) 947.
- [15] M. Suzuki, Mod. Phys. Lett. **A5** (1990) 1205; A. Hasenfratz, P. Hasenfratz, K. Jansen, J. Kuti and Y. Shen, Nucl. Phys. **B365** (1991) 79. They claim that NJL model has no predictive power at low energies because it has extra degrees of freedom of parameters due to higher dimensional interactions allowed by symmetry. The physics, however, is not determined solely by symmetry, but the precise differences in the Lagrangian reflect some differences in the fundamental physics. Each case leads to different compositeness condition, and different predictions. We can use the freedom to avoid the instability of the potential.
- [16] K. Akama, Prog. Theor. Phys. **64** (1980) 1494; **84** (1990) 1212.
- [17] J. D. Bjorken, Ann. Phys. **24** (1963) 174.
- [18] A. D. Sakharov, Dokl. Akad. Nauk SSSR **177** (1967) 70 [Sov. Phys. Dokl. **12** (1968) 1040]; K. Akama, Y. Chikashige, T. Matsuki and H. Terazawa, Prog. Theor. Phys. **60** (1978) 868; K. Akama, Prog. Theor. Phys. **60** (1978) 1900.


FIGURES

FIG. 1. The diagrams which are leading ((a), (b)) and next-to-leading ((c)–(g)) in $1/N$ and leading in $1/I$. The solid and dashed lines indicate ψ and ϕ propagators, respectively. The big-dotted line indicates the ϕ propagator with a number of one- ψ -loop insertions. The blob indicates a planar or a crossed one- ψ -loop insertion at the vertex.




(a)


(c)




(b)




(d)



(e)



(f)



(g)